

Advil gel

This information is a summary only. It does not contain all information about this medicine. If you would like more information about the medicine you are taking, check with your doctor or other health care provider. No rights can be derived from the information provided in this medicine leaflet.

Advil Gel

Read all of this leaflet carefully because it contains important information for you. This medicine is available without prescription to treat minor conditions. However, you still need to use it carefully to get the best results from it.

- Keep this leaflet, you may need to read it again
- Ask your pharmacist if you need more information or advice
- You must contact a pharmacist or doctor if your symptoms worsen or do not improve after 14 days

What this medicine is for

This medicine contains Ibuprofen which belongs to a group of medicines called non-steroidal anti-inflammatory drugs (NSAIDs), which act to relieve pain and reduce inflammation (swelling).

It can be used to relieve mild to moderate pain and inflammation (swelling) caused by backache, rheumatic and muscular pain, sprains, strains and sports injuries.

Before you use this medicine

This medicine can be used by adults and children of 14 years and over. However, some people should not use this medicine or should seek the advice of their pharmacist or doctor first.

Do not use:

- If you are allergic to ibuprofen or any other ingredients of the product
- If you have asthma, allergic runny nose (rhinitis) or itchy, red, raised skin rash (urticaria) and have ever had a bad reaction to aspirin, ibuprofen or other NSAIDs in the past
- On broken, damaged, diseased or infected skin
- If you are under 14 years of age, unless your doctor tells you to
- If you are pregnant or breastfeeding, unless your doctor tells you to!

Talk to your pharmacist or doctor:

- If you have a stomach ulcer or an ulcer in the intestine, or have had one in the past
- If you have kidney problems
- If you have asthma, or a history of asthma
- If you develop a rash when using the gel – and do not apply any more gel

Other important information

This gel may make your skin more sensitive to sunlight. Keep treated areas out of direct sunlight, but do not cover with plasters or dressings.

If you use other medicines

Before you use this gel, make sure that you tell your pharmacist about ANY other medicines you might be using at the same time, particularly the following:

- Aspirin, or other non-steroidal anti-inflammatory pain killers (if you use these medicines with Ibuprofen gel you may notice more side effects)
- Medicines to treat high blood pressure, or medicines to thin the blood (Ibuprofen gel may rarely affect these medicines)

If you are unsure about interactions with any other medicines, talk to your pharmacist. This includes medicines prescribed by your doctor and medicine you have bought for yourself, including herbal and homeopathic remedies.

How to use this medicine

Check the tube seal is not broken before first use. If it is, do not use the gel.

Pierce tube seal with the point in the end of cap.

Age	How much to use	How often to use
Adults and children of 14 years and over	4 – 10 cm of gel (50 – 125 mg of ibuprofen)	Every 4 hours Don't use more than 4 times in 24 hours.

Gently massage the gel into the affected area until it is absorbed.

Wash your hands as soon as you have used the gel.

Apply only to the skin. Avoid contact with eyes, mouth, lips or nose, or inflamed or broken skin. Do not apply plasters or dressings over the gel.

Do not use on children under 14 years, unless your doctor tells you to.

Do not use more than the amount recommended above.

Stop using the gel if a rash develops.

If symptoms worsen, or do not go away after 14 days talk to your doctor.

Possible side effects

Most people will not have problems, but some may get some.

! If you get any of these serious side effects, stop using the gel. See a doctor at once:

- Unexplained wheezing (asthma), worsening of existing asthma, difficulty in breathing, swelling of the face, tongue, neck or throat (severe allergic reactions)
- Allergic skin reactions such as itchy, red, raised rash (which can sometimes be severe and include peeling, blistering, weeping and lesions of the skin) These other effects are less serious. If they bother you talk to a pharmacist:
- Stomach pain, heartburn
- Kidney problems (particularly in people with a history of kidney problems)

If any side effect becomes severe, or you notice any side effect not listed here, please tell your pharmacist or doctor.

How to store this medicine

Store below 25°C.

Store in the original package.

Keep this medicine in a safe place out of the sight and reach of children, preferably in a locked cupboard.